copied for CAS 11:10:1946

Sta. No. H45-26

Locality: California. Kelp beds, off La Jolla

Approximately

Lat. 32 ° 50' 00" N.; Long. 117 ° 17' 15" W. (USC & GS chart 5106, 1917.)

Water: clear, salt

Vegetation: beside kelp bed.

Bottom: Temp.:

Shore: Current:

Distance from shore: about 0.3 mile Tide:

Depth of capture: bottom Depth of water: est. 5 fath.

Method of capture: three commercial fishermen — lobster trap

Collected by: Carl L. Hubbs Date: III : 3 : 1945

Orig. preserv.: form. Time:

35. Paralabrax sp.: small catch.

47A Embiotoca jacksoni → 1 (206) : 1 eg. ad. purplish with silver-blue bars on lateral scales anteriorly.

SIO

53. Sebastodes auriculatus : 1 ad. ♂,

CAS 19334 D. XIII, 13; A. III, 7; C. 14; P. 18-18 (i, vii, 10). Color similar to H45-25 (Copied for drawings 53)

Sebastodes carnatus : 1 ♀ turgid with eyed embryos. Dark areas dirty yellowish; light areas pinkish cream, including outer part of thickened pectoral rays. Light saddles along back verging toward purplish rather than cream. Fins largely yellowish, lower stripe on head golden, lower part of head bluish gray. Virtually no spotting on lower anterior surfaces.

CAS 19331

(over)

Sebastodes serriceps : 2 hl. g. - ad (all branched)

D. XIII,14; A. III, 6, P_1 --18, P_2 I, 5-I, 5, C. 14
X, III, 15 III, 6 P_1 17–17 (i, vii, 9); P_2 I, 5-I, 5; C. 14

CAS 19391

Ground color dirty yellow with blue-black bands. Pink about lips and over lower side of head to same extent on pelvic and anal spines and adjacent parts of body and fins.

(Counts transferred to Rockfish file)