

Copied for
CAS

Sta. No. H 45-27

Locality: California: outside kelp beds NW of
La Jolla Point

Lat. 32° 51' ca. 20' N.; Long. 117° 16' ca. 45' W. (USC & GS chart 5106, 1917)

Water: clear; salt

Vegetation: beyond kelp

Bottom: Temp.:

Shore: Current:

Distance from shore: Tide:

Depth of capture: bottom Depth of water: ab. 85 ft

Method of capture: wire lobster trap; brought in in water in boat.

Collected by: fishermen^(Mike?) for Carl L. Hubbs Date: III; 6; 1945

Orig. preserv.: Time:

Neoclinus blanchardi [change VII: 2: 1949]

61. *Pterognathus satoricus*: 1 subad., to

aquarium - dull chocolate, with a
little reddish glow; broken near back
by paler green blotches, less
conspicuous than green-cream blotches
on 24. D with pinkish splashes also,
on a light brown base; edge of
fin bluish white. Anal splashed
with greenish, pink and brownish;
becoming blackish within bright cream
border. Ocelli on 24, largely
metallic blue, with becoming black
just within the narrow bright
golden & yellow ring. Maxillary
pale pinkish brown, with lower &
posterior border of bright yellow;
yellow also extending into mouth
on

CAS.
To 18425

very conspicuous when mouth is open.

Central part of mouth pinkish.


Main supracocular area sooty but
posterior cuni. & nasal flaps light
tan. C. dirty greenish, with edge
broken into darker of greenish and
reddish. P, largely brown, with
bluish blotches near base.
photographed by Paul Williams

Next day was light tan with
deep brown cross bars!

Second day had almost returned
to original color - rephotographed
by Williams.

Note III: 17: has varied back & forth.

Automatically lie in a ~~po~~ sharp

arc (). A light spot
behind & below eye has become
conspicuous (white with dark border).