Copied for CAS I:16:1946

Sta. No. 1945-47

Locality: California: off La Jolla Bay, in ~~deep~~ channel submarine canyon

about

Lat. 32° 52' 20" N.; Long. 117° 16' 40" W. (USC & GS chart 5101, 1933)

Water: clear, salt

Vegetation:

Bottom: Temp.:

Shore: Current:

Distance from shore: Tide:

Depth of capture: on bottom Depth of water: "800 or 900 ft."

Method of capture: hook + line by ~~Foxy Blanks~~ Mike)

Collected by: Carl L. Hubbs Date: IV:6:1945

Orig. preserv.: 10% form + boras Time:

5. ~~Squalus~~ Squalus suckleyi ~~acanthias~~: 1 ad. ♂, unspotted - not saved.

53. Sebastodes ~~paucispinis~~: 2 ad. gr. saved. about 10 more caught. Fisherman says definitely fish were caught on bottom at same depth as goodei. Upper parts dusky olive, mid sides of smaller specimen with coarse reticulations of rich brown and bluish gray. Lower parts with a wash of orange. Upper sides with blackish brown bordered spots. ~~mandibular region rather strongly orange. Throat~~ Anal and lower part of C. ochraceous with blackish membranes. Paired fins somewhat more orange.

CAS not rec'd by CAS Mar. 1946

Sebastodes goodei: 1 subad. preserved, a larger specimen taken by Wenham. Fishermen says this is the real chilipepper. Rather strong suffusion of red, brightest in streak along lateral line. Branchiostegal region, breast and

CAS not rec'd by CAS March 1946

H45-47

belly largely white, mandibular region strongly washed with red.

Sebastodes elongatus: 1 subord. Upper 2/3 of body watery gray with some blue ventrally and considerable red toward D. base. Boldly marked with brassy green with considerable black. Breast and lower side of head largely white but other lower parts pink. Considerable red above jaws and on middle of face. First D. largely greenish, with a subbasal bluish band and pinkish tip. Second D. rather similar but with more pink. Upper & lower edges of C. pink. Rest of fin mixed pale pink and green. Pelvic and Anal pale pink with whitish edgings. Pectoral pale orange.

CAS 1845-3