

copied for CAS

Sta. No. 1445-82

Locality: Baja California: south side of
Punta Banda near base - (Gelidium area)

Lat. 31° 42' 50" N.; Long. 116° 41' 35" W. (USC & GS chart 1149, 1922.)

Water: clear, salt

Vegetation: between kelp and shore (about 75 ft.); eel grass
+ algae on bottom

Bottom: rock

Temp.:

Shore: rocky - cliffs (mostly steep)

Current:

Distance from shore: close

Tide: incoming

Depth of capture: ~~low~~ near bottom

Depth of water: 10-20 ft.

Method of capture: hook and line

Collected by: Carl D. Johnson and party Date: VI: 8: 1945

Orig. preserv.: 10% form. + borax Time: 8-11 A.M.

Discarded 2 or 3 large "black perch"

5 "kelp bass"

3 large reddish "rock cod"

2 sheepshead ♂ + ♀

♂ about 6 lbs.

♀ " 1 1/2 lb.

CAS 35. *Paralabrax clathratus*: 1 ad, 185
19360

500 47. *Embiotoca jacksoni*
sp. 1 ad (156)

CAS { 53 *Sebastes auriculatus*: 1 ad. counts taken
19376 Remains of much digested *Sibbousia*
in stomach. 212 mm.
19374 *Sebastes atrovirens*: 1 ad ♂ counts
19368 " *chrysomelas*: 2 ad, 213-214 mm.

CAS 61. *Heterostichus rostratus*: 1 ad, 213 mm
19364 *Sibbousia elegans*: 2 spec. 2(93-115)
500