

Sta. No. 145-123

Locality: California; at end of Scripps Pier, near La Jolla, San Diego Co.

Lat. 32° 52' 2" N.; Long. 117° 15' 24" W. (USC & GS chart 4809, 1928) <sup>Sheet</sup>

Water: clear salt

Vegetation: a little floating kelp

Bottom: Temp.:

Shore: Current:

Distance from shore: Tide: 5.7' at 11:21 P.M.

Depth of capture: Depth of water:

Method of capture: dip net and light

Collected by: Sach. H. Hubbs + party Date: VII: 27: 1945

Orig. preserv.: 10% + borax Time: 10:12 P.M. <sup>midnight</sup>

In addition to fishes, one large heteronereis, 2 small nereids, a few schizozoids and 3 isozoids <sup>were</sup> caught and turned over to Martin Johnson.

All Engraulis mordax mordax

8. Sardinops caerulea: 2 post-larval  
21 St. L., 23 Tot. L. } ~~25~~  
28 St. L., 31 Tot. L. } To Stanford

Tylosurus

23. Strongylura exilis: 3 spec.

183, 282, 434 mm.

To Stanford. SU 40757

Cyprinocentrus californicus: 1 ad.,  
314 mm.

To Stanford SU 40772

40.A - Leuresthes tenuis: 42 post larval (7-13.5)  
(lengths taken) at S.D.O. (8-10mm.)?

TO Stanford { 137 yg. (lengths taken) + 4 ad.,  
126, 141, 142, 149 mm. st. length.

at S.D.O.  
40 { Atherinops affinis: cedrosensis x  
littoralis: 1 adult (disc.)  
Atherinops affinis cedrosensis:  
series of adults. (discard)

Atherinopsis c. californiensis:  
101 spec. 45-168 mm. lengths taken.  
~~spec.~~  
To CAS  
~~To Stanford.~~

61 Hypsoblennius gilberti 4(19-20)

Leuresthes tenuis

Observed spawning VII:26 by  
Bob Barry

Observed a few spawning when coming  
off the pier at midnight VII:27

None observed by Gordon Tucker on  
VII:28.