

Sta. No. 445-188

Locality: California: off Coronado Strand

Lat. ° 'N.; Long. ° 'W. (USC & GS chart, 19...)

Water:

Vegetation:

Bottom: sand, mud and shell

Temp.:

Shore:

Current:

Distance from shore:

Tide:

Depth of capture: 24-25 fath

Depth of water: 34-25 fath.

Method of capture: Calif. Acad. pipe dredge

Collected by: Carl H. Hubbs and E. W. Scripps

Date: XI: 8: 1945

Orig. preserv.: 10% form. + borax

Time:

6A Raja inornata? : 3 egg cases

55A Uchelus quadriseriatus : 4 ad. 1 (58.5-60.5)

Color note, XI: 10 : Both sexes with rich red-brown spots just below lateral line. Males with deep lemon-orange throat patch and blackish branchiostegals. ♀s with only a trace of this color.

Counts in group 55 file

56A Odontaspis trispinosa : 1 hl. yr. (46.5)

Count in group 56 file

64C Citharichthys xanthostigmus : 1 ^{sub} ad. (116.5)

Count in group 64 file

64D Citharichthys stigmaeus : 27 yg.-ad.

Counts in group 64 file: 23 (30.5-85.5)

(Only 23 found 8 Oct. '62 JJC)

(over)

64. ^B Pleuronichthys verticalis : 2 yg. 35-41 mm.

Counts in group 64 file.

Gill rakers 7 (with one extra interpolated on one side of one)

Interorbital spine moderately well-developed, much more so than in ritteri of like size. Colors not bright (thus unlike ritteri yg.). Rather numerous small to large blotches some with dark centers or dark edges and ~~more~~ a fine brown network on a light ground color near the D. and A. fins. D. and A. boldly streaked with black and white. The basal part of the C. is largely black. Pectoral jet black.

64 A Symphurus atricaudus : 10 ad.

8(105-143)

(Only 8 specimens found 24 Sept. '62 JFC)