

Sta. No. H 45-189

Locality: California: Off Coronado

Lat.° 'N.; Long.° 'W. (USC & GS chart, 19.....)

Water:

Vegetation:

Bottom:

Temp.:

Shore:

Current:

Distance from shore:

Tide:

Depth of capture: 46-55 fath

Depth of water: 55 to 50 to 46 fath.

Method of capture: Calif. Acad. pipe dredge

Collected by: Carl H. Hubbs on E. W. Scripps Date: XI: 8: 1945

Orig. preserv.: 10% form. + borax

Time:

Icelinus

(2 ad) 74-84 mm

55. Tarandachthys tenuis: 1 ad. ♂, with

lower part of head sooty and pelvics and anal sooty to blackish (a point not noted by Bolin); and one ♀ with

lower parts and fins ~~to~~ white. In both sexes the specks along the lateral

line plates are bright coppery and there is considerable ^{similar} color in the

anterior part of the row of spots below the lateral line. There is also

considerable golden yellow specks and reticulations on the anterior back. Counts in group 55 file.

60. Poecilichthys notatus: 5 bl.-gr. (102-116)
Counts in group 60 file.

64A Microstomus pacificus: 1 (74)
An. Young completely transformed
Counts in group 64 file

64B Symphurus atricaudus: 13 adults ~~(90-137)~~ (90-137)

12 L; 1 R. (entered in group 64 file)

All 13 have liver on right side and coils
of intestine on left.

1 L. and 1 R. photographed

Dorsal specimen 90 mm.

Ventral, 96-137 mm.