Kelpfish Copied for CAS VIII:30:1947 Sta. No. H 46-25

Calif:
Locality: east end Catalina Island near Church Rock

Lat. 33 ° 18.5 'N.; Long. 118 ° 23 'W. (USC & GS chart 5101, 1933)
Water:
Vegetation:

Bottom: Rock Temp.: 57.5
Shore: Current:
Distance from shore: 200 yds Tide:
Depth of capture: 10 fm Depth of water: 10 FM
Method of capture: baited line
Collected by: Conner for Hubbs and J. W. Schipps Date: Feb 8, 1946
Orig. preserv.: 10% form + borax Time:

61. Neoclinus uninotatus Hubbs : 1 ad. ~~blanchardi~~

new sp. as of VII:2:1949 — Clerd/Hubbs. Iden. changed & Book V:11:1954. Folbtt ratified - S.L.H.

Color note taken at 330 II : 9

CAS

Maxillary pocket blackish green with bright chalky yellow border; the upper border suffused with dusky, the lower clear. Body brownish pink with deep purple bars, becoming purple-gray below. The region about upper base of pectoral deep red. Some small dapples of red on the D. fin above the dark bars of the body. A blackish spot between 1st 2 dorsal spines with a red tinge and with a yellow border. The barest trace of another non-ocellated spot between 7th and 10th spines. Most of spinous D. dusky with a reddish wash. Soft D. and C. rays with red spots near base and orange spots

near tip. Anal fin mostly purplish with a little dusky red near base and a very few orange spots near tip of rays. Upper pectoral rays tipped with yellow. Pelvics blackish with a little red. Head mostly dull purplish red, with a little dusky greenish on upper part of membrane between branchiostegal membranes. Premaxillary and maxillary deep pink.