Copied for CAS VIII.5.1947

Sta. No. H46-112

Locality: California: off La Jolla Pt. (about 2½ miles W.)

Lat. ° '.N.; Long. ° 'W. (USC & GS chart , 19)

Water:

Vegetation:

Bottom: Temp.:

Shore: Current:

Distance from shore: Tide:

Depth of capture: Depth of water: ab. 350 ft.

Method of capture: set lines

Collected by: agent fisherman, brought by Wm. Wareham Date: X:27:1946

Orig. preserv.: form. + borax Time:

61. Chilara taylori 1 (116 13 mm)

CAS

53. Sebastodes ovalis Ayres ?: 1 ripe ♀ 1 expelling minute embryos. P_{11} P_2 A. salmon, dusties on membranes. C. dusky salmon with blackish eyes. D's olive salmon with numerous green-black specks. numerous fine blackish specks wrapper ⅔ of body. Body tan with a weak orange tinge orange-white on belly and throat: a little pink on lower part of head

counts in group 53

Gtoglididiceum taylori : regurgitated by rock cod, partially digested. 1 lb. 9 oz. (116)

To Stanford XII:2:49

Sebastodes differs from description in Jordan and Evermann — 16 D_2 instead of 14, 9 A_2 rays instead of 8. Sc and 9. spine but little longer than 3rd, not quite reaching tip of 3rd when depressed, only about 1/10 longer and contained about 2.6 in head. Peritoneum dusky purple not black. Certainly looks like Fig. 100 in Woolford's Handbook