Sta. No. H46-224 (1542-46)

Locality: Revilla Gigedo Is.: Grayson Cove on W shore of Socorro Id.

Lat. 18° 44.4' N.; Long. 110° 59.6' W. (~~USC & GS~~ HO chart 1687, 1925)

Water: clear, blue

Vegetation: ~~at no~~ little Considerable coral, especially on submerged rock reefs

Bottom: mostly boulders and reefs. Temp.:

Shore: 2 coral beaches with reef bottom Current: little

Distance from shore: close and cliffs at side Tide: moderately low

Depth of capture: Depth of water: to 5 fath.

Method of capture: observation (except a few mollusks)

Collected by: Zaca Exped. Date: VIII: 26: 1946

Orig. preserv.: mollusks dry Time: midday & early P.M.

A few mollusc shell picked up. Gastropods + large chitons on rocks above water.

Search for water mentioned in Sailing Direction proved fruitless. Either spring is no longer in existence or it is not at point described.

Fishes seen: most of common reef fish seen at other point.

Also — see over

Strongylura

23. ~~Tylosurus~~ sp. One in channel against shore, in strong surge. Looked like T. stolzmanni.

40. Mugil sp. young in channel.

Cirrhitus rivulatus (?)

52. Cirrhitid (?) about 10" long, was under an under-cut rock; on my approach it onshore it swam toward me, until it was temporarily stranded. Some small fish were with it & possibly being guarded.

59, 61 Bathygobius, blennies & a considerable variety of other small fishes, including yg. of coral reef pomacentrids, etc., in the pools on the flat reef on 2 levels, both above a submerged ledge rich in coral. good organism collecting station.

MAR 3 - 1954

Inf. to Bayard Brattstrom for publication

The towhee (?) with chestnut on ~~[illegible]~~ flanks sings prominently and often. One was flushed from a nest in ~~thigh~~ ~~[illegible]~~ bushes near coast. Eggs with rather indistinct dark markings on bluish-gray base.