

To UCLA V: 12:53

Sta. No. H46-226 (1544-46)

Locality: Revilla Gigedo Islands:
at anchorage of Binnero Cove, near
S end of W. shore of Pocosco Island

Lat. 18° 43.1' N.; Long. 110° 58.0' W. (USC & GS chart 1687, 1925)

Water: clear

Vegetation: none

Bottom: Temp.:

Shore: rocky Current: little

Distance from shore: 1/8 - 1/4 mile Tide:

Depth of capture: surface Depth of water: 13 fath.

Method of capture: dip-nets

Collected by: Zaca E. ... (Hutch-Flynn VIII: 26: 1946)

Orig. preserv.: 1090 fresh + frozen Time: 8:30:11:50 P.M.

In striking contrast to H46-220 collection, invertebrates were scarce.

5 Carcharias platyrhynchus.
bl. gy. - ad. rather common -
fed on wounded flying fish
none caught.

23. Tylosurus stolzmanni:
~~Stenogobius~~
ser. of bl. gy.

Cyprinus -
series of 2 mm sp., with
red Pectorals, also caught
at H46-220 and seen
around island.

over.

Anter edge of pectoral bluish to
 main part water, red; ~~at~~
 inner edge whitish. Lower
 side of pectoral with
 red largely concealed, except
 distally, by blue-silver
 along the fringe-like rays.
 A trace of pink on upper
 side.

Cyprichromis callostrum : 1 ad.

Cyprichromis _____ : 2-3
 ads - large sp. with
 most of D black.

40. ^{Atherinella} Euryostole ericrha : ⁵³⁽³⁴⁻⁵⁴⁾ ~~53~~,
 bl. eye - od. ; at var. surface.

40A Chaenon Muzil proboscideus 5 yf. 6(35.5-38)
 4(24-25) 5 " Therichthys 75

33A ~~42~~ Acanthurus triosteus : 7
 4 ~~postlarva~~ yf. with trenchant belly.

34. Tetraodon 1 yf. - steindachneri
 61A Ophiochromis sp. ~~10(20-51)~~
seri 7 yf. 1 ad.

34A Canthipaster punctatissima 1(15.5)