Sta. No. H47-89

Locality: Calif: Santa Cruz I, north shore of Fraser pt. Tide pool 8 × 25'

Lat. 34 ° 03.65 'N.; Long. 119 ° 55.52 'W. (USC & GS chart 5115, 1945)

Water: Clear salt

Vegetation: Brown and coralline algae, very scarce

Bottom: Rocky Temp.:

Shore: ~~Tide pool~~ - Rocky Current:

Distance from shore: Tide pool Tide: low F.0.9 at 11 (PM)

Depth of capture: 2 ft max Depth of water: to 2 ft.

Method of capture: Poison - derris (moderately thorough). "E.W. Scripps"

Collected by: Carl L. Hubbs & party Date: 28-IV-47

Orig. preserv.: 10% formalin Time: 1200 hrs

Pool on top of flat with reef, with rocky projection in center but rather poor coralline development.

All fish saved.

Pool about 1 foot above low tide of day.

55. Artedius lateralis: 1 ad.

Oligocottus ~~Leptocottus~~ rubellio: 19 ad., ♂1 + 0, 35-55 mm. 19(35-55)

Oligocottus ~~Leptocottus~~ snyderi: 2 ad. 2(50-50)

Clinocottus analis: 4 ad. 4(71-84)

Monterycya recalva: 3 ad. (covers)

61. Sibbonsia metzi : 1 lbl. gr.

Sibbonsia sp. : 1 postlarva,
4 juvenil, 6 lbl. gr. ad.

Epigeichthys atropurpureus : 1 yy.

500 fish. yy. coll.

61 Xererpes fucorum : 4 ad. 3 (107-117)