

Copied from  
CAS VIII:5: (1947)

Sta. No. H47-136

Locality: Calif. Santa Rosa I. Becher's  
Bay, Vale Ranch. Pool ground and  
hook and line used in one pool near pier

Lat. 34° 00.50' N.; Long. 120° 02.78' W. (USC & GS chart 5115, 1945)

Water: clear, salt

Vegetation:

Bottom:

Temp.:

Shore:

Current:

Distance from shore:

Tide:

Depth of capture:

Depth of water:

Method of capture: W.M. hook + line and black beef 40' garrison

Collected by: Mrs. Lynch (= Helen S. Lynch) Date: Rec'd 5 July 47

Orig. preserv.: 10% formalin. Time: May-June, 1947

~~Tide pool collection~~

*Embiotoca lateralis*

1 (206)

47A) *Taeniotoxa lateralis*: 4 ad 3 (133-217)

hook + 47B) *Embiotoca yuhsani* → 7 (119-247)  
line → 9 (55-247) → 2 (55-114)

47C) *Phanerodon furcatus*: 2 ad. (163-176)

*Rhacochilus*  
47D) *Damalichthys vacca*: 1 ad ♀ with  
6 yg. 7 (50-279)

47) *Gnathypops aurora*: 4 ad (79-110)  
*Micrometrus*  
Probably tide pool poisoning

55) *Clinocottus analis australis*: 11 ad  
11 (70-100)

Tide pool poisoning 55) *Corte dius lateralis*: 1 ad (89)

55) *Oligocottus*  
~~*Damalichthys*~~ *myleri*: 5 unad  
5 (35-57)

(over)



61A Cebidichthys violaceus : 1 large ad (314)

"High and dry in rocks at low tide"

6b Xerocypus bicolor : 1 ad. (114)

Sibbomsia metzi : 3 yg. - 6l. gr.

Tide pool  
pursuing

CAS