Sta. No. H48-177

Locality: California: near Santa Barbara. ½ mi. off Hendrys Beach just about 1 mi. up the coast from Santa Barbara

Lat. ° 'N.; Long. ° 'W. (USC & GS chart , 19)

Water:

Vegetation: ab. 1/4 mi. outside of kelp

Bottom: Temp.:

Shore: Current:

Distance from shore: ab. ½ mile Tide:

Depth of capture: Depth of water:

Method of capture: blasting

Collected by: Elmer R. Noble Date: VI: 15: 1948

Orig. preserv.: ~~formalin~~ Time:

Not all fish saved.

46. Sebastomus serriatus: ~~not many~~ killed, not preserved

47 A. Hypsurus caryi: 6 ♂ (4 taken to SIO) → 2 (155-162)

X-rayed by C.L. Turner

Brachyistius frenatus: 7 ad. (4 to SIO) 4 (87-125)

47 Phanerodon furcatus: 3 ad. (1 to SIO) 1 (125)

Embiotoca jacksoni: 1 ad taken to SIO with thick lips; not taken

(1 large Embiotoca) 1 (161)

48. Chromis punctipinnis: 1 ad.

53. Sebastodes paucispinis: 37 yg.

yg. Killed by the 1000's.

Sebastodes vexillaris: 2 hl. gr. → 1 (180) R.J.K. Feb 59 (1 to SIO)

1 large Sebastodes discarded in field looked like vexillaris

60. Porichthys myriaster: 1 hl. gr.

" notatus: 2 "

61. Heterostichus rostratus: 1 hl. gr.