Sta. No. H 48–258

Locality: Baja California: vic. Punta China, mi. S. of mouth of Arroyo Santo Tómas

Lat. ° 'N.; Long. ° 'W. (USC & GS chart , 19)

Water: clear, salt

Vegetation: about kelp beds (Macrocystis).

Bottom: Temp.:

Shore: rocky Current:

Distance from shore: Tide:

Depth of capture: Depth of water:

Method of capture: hook (Mex. fishermen for lobster bait)

Collected by: Carl L. Hubbs Date: IX: 4-5: 1948

Orig. preserv.: 10% form. Time:

Picked up IX:5 — apparently caught over last 2 days.

50. Pimelometopon pulchrum: ser. hl. gr. — 2d. (not saved).

53. Scorpaena guttata: 1 ad.

Sebastodes ~~rosaceus~~ (miniatus): ser. hl. gr. — 2d. (none saved).

53B Sebastodes ac roseus, n. sp. 1 ad. (161 mm.) saved.
Typical color: strong purple border to pink spots.

53A Sebastodes constellatus: 1 hl. gr. (saved) + 1 2d. :.....1(187)

Sebastodes vexillaris: ser. hl. gr. — 2d. (none saved).

* Sebastodes carnatus: about 30 (169-184 mm.) hl. gr. – 2d., typical (2 saved).

53C Sebastodes rubrivinctus: 1 hl. gr. (162) (saved).

* 1(184) gift to Zool. Inst., Leningrad

(over)

Fish Data Sheet–1m-8,'46 (9814s) 97

H48-258 (cont)

55. Scorpaenichthys marmoratus:
1 ad., typical oliviaceous inshore type, not saved.

60. Caulolatilus princeps anomalus:
several ad. (not saved).