

State or Country: CALIFORNIA

Field No. H 52-79

County: Humboldt

Map:

1

Locality: Trinidad Head Near pier

Lat.

N., Long.

W.

Water: Murky salt. Visibility 4 feet

Vegetation: Egregia, Nereocystis

Bottom: Rocky

Cover: Egregia + rocks

Temp: About 10°C

Shore: Rocks + sand

Current: EAST

Dist. offshore: 30 feet

Stream width:

Depth of capture: 0-18 feet

Depth of water: 0-18 feet

Collected by C. Limbaugh + A. Rehnitz

Tide: In coming

Date: VI: 21, 1952

Method of capture: Rotenone + diving

Orig. preserv.: 10% FORMALIN

Time: 1000-1300

Embiotoca47 *Taeniotoca lateralis* (Agassiz) 3 yg. 56-60 mm.

SL.H. 1849
 IV: 29
 1960
 v: 3:60
 (53A) *Sebastes melanops* (Gir.) 34-44 39-48 Counts
 (B) *Sebastes mystinus* (Jordan & Sill.) 1 yg. 42 mm. Counts
 (C) *Sebastes pinniger* (Jordan & Sill.) 1 yg. 34-41 mm. Counts
 (D) *Sebastes verillaris* (Jordan & Sill.) 10 yg. 21-32 mm. Counts
 For *S. caurinus* (Richardson)

54 *Hexagrammos decagrammus* (Pallas) 20 yg. ad 69-215 mm
Hexagrammos 27 (69-215)

55 *Artedius harringtoni* (Starks) 7 ad: 25-62 mm.*Ascelichthys rhodurus* Jordan + Gilbert 3 ad: 82-105 mm.*Enophrys bison* (Girard) 1 ad: 233 mm.55 *Hemilepidotus spinesus* (Ayres) 20 yg. ad: 28-148 mm.*Nautichthys oculofasciatus* (Girard) 1 ad: 105 mm.*Scorpaenichthys marmoratus* Girard 1 yg: 50 mm.56 *Liparis flerox* (Jordan + Starks) 1 ad: 69 mm.56 *Liparis lucensis* Ayres; 2 yg. ad 24-29 mm (24.5-75.5)60A *Gobiesox* sp. 9 (9-10)61 *Apodichthys flavidus* Girard 2 yg (33-42 mm)61 *Anarrichthys ocellatus* Ayres 1 yg: 51 mm 1 (50)61C *Chirolophis nigrator* (Jordan + Williams) 5 yg. ad (31-104)61 *Xiphiaster* ~~*Scorpaenichthys*~~ *atropurpureus* (Kittlitz) 2 ad: (190-252 mm)61E *Pholis schultzi* 5 (78.5-106)