

State or Country:.....Field No. **SIO60-285**

County:.....Map:....., 1.....

Locality: **Eastern North Pacific Ocean**Lat **29° 58.0'** N., Long. **128° 48.2'** W.Water: **to 30° 05.8' N., 128° 18.5' W.**

Vegetation:

Bottom:

Cover:Temp.:

Shore:Current:

Dist. offshore:Stream width:

Depth of capture: **500m ± 55°** Depth of water: **2400 + fms**Collected by **Tethys expedition - SIO**Tide:Date: **14 Aug 1960**Method of capture: **10 ft. midwater trawl**Orig. preserv.: **10% form** Time: **1600 - 2345****Tethys sta # 32****65A Lophodolus diadema indicus 1 (72.5)****35A Howellia ~~pammelas~~ ? n. sp. #1 1 (22.5)****10A Idiacanthus antrostomus 1 (92)****10B Sternoptyx diaphana 1 (23.1)****10B Sternoptyx diaphana 5 (7.6-17.2)****** or SIO 60-278****10C Argyropelecus lynchus lynchus 1 (14.6)*****10D Argyropelecus pacificus 1 (27.5)****10E Gonostoma atlanticum 1 (46.7)****25A Diogenichthys atlanticus 1 (17.5)****25B Ceratoscopelus townsendi 1 (21.9)****25C Ceratoscopelus sp. 3 (20.3-21.5)****10F Chauliodus sloani 1 (56.4)****10G Gonostoma ebelingi 1 (46.4) + 1 head****30A Stylephorus chordatus 1 (118)****25D Lampadena urophos 1 (24)****25E Tazaraichthys bathyphilus 1 (83)****31A Poromitra oscitans ~~parrifres~~ 3 (54-79)****31B Scopelogadus mizolepis bispinosus 1 (54)****~~25 Lampanycetus niger 3 (58-71)~~****25 Lampanycetus steinbecki ? 2 (29-58)****25 Notolychnus valdiviae 3 (21-23)****25 Triphoturus nigrescens ? 1 (31)****25 Scopelarchus analis 1 (76.0)****25 Tazaraichthys paurolychnus 1 (83)****31 Scopeloberyx opisthopterus 2 (20-34)**

- | | | | |
|----|---------------|---------------|-----------|
| 25 | Nannobrachium | bristori | 1(73) |
| 25 | Nannobrachium | hawaiiense | 2(59-65) |
| 10 | Ichthyococcus | irregularis | 1(14) |
| 26 | Serrivomer | sp. (damaged) | 1(315) |
| 26 | Cyema | atrum | 1(90) |
| 23 | Exocoetus | sp. | 1(11) |
| 10 | Cyclothone | pseudopallida | 7(19-35) |
| 10 | Cyclothone | signata | 19(22-30) |
| 10 | Cyclothone | acclinidens | 4(27-47) |