

State or Country: \_\_\_\_\_ Field No. **SI061-636**  
County: \_\_\_\_\_ Map: \_\_\_\_\_, 1  
Locality: **South China Sea**

Lat. **10° 35.1'** N., Long. **112° 51.0' E.**

Water: \_\_\_\_\_  
Vegetation: \_\_\_\_\_

Bottom: \_\_\_\_\_  
Cover: \_\_\_\_\_ Temp.: **28.7°C. surface**

Shore: \_\_\_\_\_ Current: \_\_\_\_\_  
Dist. offshore: **1500 mwo** Stream width: \_\_\_\_\_

Depth of capture: **0-ca. 800 m.** Depth of water: **2651-2378 m.**

Collected by **T. Matsui, Chaiyo, Kosol, VanLandingham**

Tide: \_\_\_\_\_ Date: **12 June 1960**

Method of capture: **2 m. stramin net**

Orig. preserv.: **10% form. + borax** Time: **1830-2055**

**Scripps** **Naga Sta. 60-512, (Cruise 5-6)**

- Triphoturus
- |               |  |                    |
|---------------|--|--------------------|
| 25            | Lampanyctus nigrescens                       | 7 (14-26)          |
| 25            | Benthosema saborbitale                       | 3 (18-24)          |
| 25            | Lampanyctus basiti                           | 2 (27-30)          |
|               | turneri                                      |                    |
| 10            | Cyclothone alba                              | 2 (                |
| 25            | Ceratoscopelus townsendi <sup>warmingi</sup> | 1 (19.3)           |
| <del>31</del> | <del>Direlmus sp. argenteus</del>            | <del>1 (6.5)</del> |
| 25            | Lampanyctus aletus                           | 12 (11-43)         |
| 32            | Forcipiger sp.                               | 1 (5.5)            |
| 25            | Hygophum sp. atratum                         | 3 (11-13)          |
| 10            | Sternoptyx obscura                           | 8 (5-10)           |
| 10            | Eustomias (Nominostomias) sp                 | 1 (8.2)            |
| 25            | Omosudis lowii                               | 1 (18)             |
| 10            | Stomias affinis                              | 1 (47)             |
| 10            | Idiacanthus fasciola                         | 5 (22-48)          |
| 25            | Taaningichthys bathyphilus                   | 1 (47)             |
| 31            | Diretmoides pauciradiatus                    | 1 (6.5)            |
| 10            | Cyclothone pseudopallida                     | 10 (21-33)         |
| 25            | Diaphus thiollieri                           | 1 (16)             |