~~State or~~ Country: Canada Field No. SI063-599

County: British Columbia Map: 1

Locality: Howe Sound: Keats Island, [redacted]: Twin Island, NW. tip

Lat. N., Long. W.

Water: salinity: 16.5‰ at surf.; 27.9‰ at 50 ft.

Vegetation: Laminaria sparse to 30 ft.; coralline algae below 30 ft.

Bottom: cliff; large rocks with sand in interstices sloping to soft sand at 50 ft.

~~Cover:~~ Temp.: 60°F. at surf.; 48°F. at 50 ft.

~~Shore:~~ ~~Current:~~ [redacted]

Dist. offshore: 100 ft. ~~Stream width:~~ thermocline at 25 ft.

Depth of capture: 20 to 50 ft. Depth of water: 20 to 50 ft.

Collected by R. Rosenblatt, D. Creed, J. Webster, D. Gordon, & G. Cowan

Tide: plus 9 ft.; falling Date: 29 May 1963

Method of capture: Rotenone, diving, dipnet

Orig. preserv.: 10% formalin Time: 1130 to 1430

[redacted] BC63-732

55A Artedius harringtoni 7(35-59)

55B ~~Artedius~~ Ruscarius meanyi 2(28-34)

55C ~~Gilbertidia~~ Psychrolutes sigalutes 1(51)

Field Data Sheet: Fish—12m-8,'60(B2277m)997