

SIO 64-56

State or Country: BAJA CALIFORNIA, MEX Field No. MAG BAY B-17

County: \_\_\_\_\_ Map: HO # 1636, 1922

Locality: SOUTH OF PTA. REDONDO

Lat. 24° 30.3' 79.8' N, Long. 111° 59.4' W

Water: — Visibility ca. 10 feet max

Vegetation: \_\_\_\_\_

Bottom: ROCKY WITH FEW SAND PATCHES

Cover: \_\_\_\_\_ Temp.: 63.6°F (TOP)

Shore: \_\_\_\_\_ Current: \_\_\_\_\_

Dist. offshore: 500 YDS Stream width: \_\_\_\_\_

Depth of capture: 60'-70' Depth of water: 70'

Collected by MAG BAY "D" B. N. Kobayashi & party

Tide: \_\_\_\_\_ Date: 3 FEB, 1964

Method of capture: CHEM. FISH POISONING

Orig. preserv.: 10% FORMALIN Time: 0840-0925 HRS

*Isolated area consisted of a rocky bottom with a few scattered sand patches. Surge was strong.*

- 35A *Rypticus bicolor* 1(137)
- 26A *Gymnothorax parameensis* 1(213)
- 26B *Muraena argus* 1(395)<sup>106</sup>
- 35B *Serranus fasciatus* 4(32-~~37~~)
- 35C *Alphesthes galapagensis* 13(36-77)
- 59 *Lythrypnus pulchellus* 2(16-17)
- \* 61 *Paraclinus magdalenae*: PARATYPES  
..... 3(22-37)
- 61 *Paraclinus magdalenae*: HOLOTYPE  
..... 1(39)
- 35 *Apogon retrosella* 9(40-56)
- 61 *Rurula azalea* 2(40-70)
- 32 *Heterostichus nigrostris* 2(50-55)
- 50 *Nicholsina denticulata* 4(37-50)
- 48 *Chromis atrilobata* 11(62-82)
- 34 *Sufflamen verres* 11(65-77)
- 61 *Heterostichus rostratus* 1(133)
- 34 *Diodon holacanthus* 2(77-80)
- 45 *Calamus brachysomus* 1(187)
- 61 *Labrisomus xanti* 4(94.3-127.5)
- 61 *Labrisomus multiporosus* 1(111)
- 48 *Eupomacentrus rectitraerum* 1(41.4)
- \* gift to USNM, July 1969

48	<i>Eupomacentrus flavilatus</i>	5(22-53)
32	<i>Johnrandallia nigrirostris</i>	2(50-55)
61	<i>Ogilbia</i> sp.	1(52)