

SIO 64-62

State or Country: Baja California, Mexico Field No. MAG BAY B-23
 County: _____ Map: HO # 1636, 1922
 Locality: SE of Puerto Cortez, Bahía Almejas
 Lat. 24° 27.3' N., Long. 111° 46.0' W.
 Water: Visibility poor - 3-5 feet max
 Vegetation: _____
 Bottom: Mud, sand, rock outcrops
 Cover: _____ Temp.: ST = BT = 62.1°F
 Shore: _____ Current: _____
 Dist. offshore: 15- yards Stream width: _____
 Depth of capture: 10-15 feet Depth of water: 15 feet
 Collected by: Mag Bay "B" B.N. Kobayashi & party
 Tide: _____ Date: 5 FEB 1964
 Method of capture: CITEM-FIST POISONING
 Orig. preserv.: 10% FORMALIN Time: 1435-1550

Area poisoned was adjacent to a rock outcrop (above surface of water) on which many birds were perched - The bottom consisted of mud and/or sand with small rocky outcrops - Rocks were 2-7 feet in diameter and up to 3 feet high - There was little surge.

- | | | |
|---------------|---|----------------------|
| 32A | <i>Pomacanthus zonipectus</i> | 1 (230) tank |
| 34A | <i>Balistes verres</i> | 1 (276) tank |
| → 35A | <i>Paralabrax maculatofasciatus</i> | 31 (35-86) |
| 35B | <i>Epinephelus analogus</i> | 1 (71) |
| 61 | <i>Paraclinus sirus</i> | 2 (25-32) |
| 35 | <i>Apogon retrosella</i> | 6 (39-53) |
| 46 | <i>Parques viola</i> | 2 (73-76) |
| 48 | <i>Abudefduf troschelii</i> | 2 (48, 2-51) |
| 59 | <i>Microgobius brevispinis</i> Ginsburg | 3 (38-44) |
| 61 | <i>Ogilbia</i> sp. | 2 (52-59) |
| 61 | <i>Ogilbia ventralis</i> | 2 (52-57) |