

State or Country: Mexico Field No. SI065-265 ✓

County: Baja Calif Sur Map: _____, 1

Locality: Isla San Jose (Gulf of Calif)

Lat. 24°52.25' N, Long. 110°37.0' W.

Water: _____

Vegetation: _____

Bottom: _____

Cover: _____ Temp.: _____

Shore: _____ Current: _____

Dist. offshore: _____ Stream width: _____

Depth of capture: _____ Depth of water: 20 ft

Collected by R. Rosenblatt + ply

Tide: _____ Date: 7 July 1965

Method of capture: cherryfish + SCUBA

Orig. preserv.: _____ Time: 1200

RR65-11 (See drawing behind SI065-260) can

| | | |
|-------|-------------------------------------|---------------|
| 32A | <i>Pomacanthus zoricpectus</i> | 1(275) tank |
| 35A | <i>Dermatolepis dermatolepis</i> | 1(307) tank |
| 35B | <i>Alphestes galapagensis</i> | 7(29-148) |
| 65A | <i>Antennarius sanguineus</i> | 1(43) |
| 35C | <i>Paralabrax auroguttatus</i> | 2(37-94) |
| 35D | <i>Rypticus bicolor</i> | 8(67-147) |
| 61A | <i>Acanthemblemaria macrospilus</i> | 5(35-43) |
| 61B | <i>Labrisomus xanti</i> | 42(21-117) |
| 50A | <i>Scarus</i> sp. | 1(13) |
| 26A | <i>Muraena lentiginosa</i> | 1(272)* |
| 26B | <i>Gymnothorax panamensis</i> | 17(70-254) |
| 31A | <i>Myripristis leiognathos</i> | 1(99) |
| 34A | <i>Canthidaster punctatissima</i> | 19(39.5-58.5) |
| 35E | <i>Apogon retrosella</i> | 52(18-80) |
| 46A | <i>Pareques viola</i> | 89(24-69) |
| 48A | <i>Pomacentrus rectifraenum</i> | 13(23-85) |
| 48B | <i>Chromis atrilobata</i> | 83(23-55) |
| 50A | <i>Halichoeres dispilus</i> | 14(19-103) |
| 53A | <i>Scorpaena mystes</i> | 4(20-109) |
| * 57A | <i>Doryrhamphus melanopleura</i> | 5(48-49) |
| 59A | <i>Coryphopterus urospilus</i> | 5(9-37) |
| 59B | <i>Lythrypnus dalli</i> | 37(10-25) |
| 59C | <i>Gobiosoma puncticulatum</i> | 3(11-22.5) |
| 59D | <i>Gobiosoma</i> sp. | 17(18.5-25) |
| 58A | <i>Gymneleotris seminuda</i> | 10(19-32) |
| 58B | <i>Chriolepis zebra</i> | 26(19-32) |
| 61C | <i>Ophioblennius steindachneri</i> | 5(44-50) |

| | | |
|-----|--|-----------------------------------|
| 61D | <i>Axoclinus carminalis</i> | 2 (26) |
| 61 | ■ <i>Enneanectes cteniceps</i> | 482 1 st ser. (9-43.5) |
| 59E | <i>Lythrypnus pulchellus</i> | 28 (10-21) |
| 32 | <i>Holacanthus passer</i> | 3 (87-170) |
| 52 | <i>Cirrhitichthys oxycephalus</i> | 10 (52-75) |
| 50 | <i>Thalassoma lucasanum</i> | 44 (4.5-88) |
| 61 | <i>Starksia spiriperis</i> | 3 (30.8-37.4) |
| 61 | <i>Protobleptaria bicirris</i> | 2 (15.5-21.5) |
| 61 | <i>Acanthobleptaria crockeri</i> | 1550 (19-41) |
| 61 | <i>Aporozia barbarae macrolepis</i> | ca. 1375 (25-53) |
| 53 | <i>Scorpaenodes xylis</i> | 143 (21-77) |
| 61 | <i>Starksia cremnobates</i> | 27 (15.5-44) |
| 61 | <i>Parastathmonotus sinuscalifornici</i> | 1 (31) |
| 61 | <i>Ogilbia</i> sp. | 21 (25-66) |

*Gift to E. Tortonese, 8/18/1972