

State or Country: Field No. SIO 67-223
 County: Map: 1
 Locality: Tanner Bank, 6 repetitive southward
1-mile-drift from buoy
at Lat. 32°41.9' N., Long. 119°07.7' W.
 Water:
 Vegetation:
 Bottom:
 Cover: Temp.:
 Shore: Current:
 Dist. offshore: Stream width:
 Depth of capture: bottom Depth of water: 16-40 fms.
 Collected by Lo-chai Chen, R. McConnaughey and party on Oconostota
 Tide: Date: 26 Oct 1967
 Method of capture: hook and line
 Orig. preserv.: Time: 0900 ~ 1600

Sebastodes rosaceus ~200 color scarlet red
 in shallower water, become somewhat
 orange in deeper water, distinctively
 different from those collected in
 La Jolla Canyon.

S. chlorostictus 3

S. constellatus ~60

S. umbrinus ~15 dark marks on back
 somewhat greenish, not evenly distributed,
 rather sparse thus differ from those
 collected in La Jolla Canyon.

S. miniatus 2 one with yellow marks
 on back, another one has no dark
 marks on back, different from any S. miniatus
 I have seen. This fish was unfortunately
 filleted by somebody.

S. paucispinis 2

S. levis 1

S. serriceps 3

S. serranoides ~3

S. elongatus 1

S. vexillaris 1

S. mystinus 2

S. jordani thrown out by S. pserranoides
 and S. rosaceus

juveniles

S. _____

S. _____

Caulolatilus princeps 50 ~ 100

Pimelometopon pulchrum ~ 15

Citharichthys sp ~ 30

on comp

Isurus glaucus

1

Prionace glauca

3

} caught at surface

Chen

53 *Sebastodes jordani*

4 (78-89)

53 *Sebastodes* sp.

2 (67-69)

53 *Sebastodes constellatus*

→ 3 (163-299)

53 *Sebastodes umbrosus*

9 (144-188)

53 *Sebastodes rosaceus*

→ 39 (134-204)

53 *sebastes hopkinsi*

1 (62)