

Trawl
50

State or Country: Central North Pacific Field No. SI068-486
 County: Between New Bank Map: _____, 1
 Locality: and Midway

Lat. 27° 56.9' N, Long. 177° 53.4' W.
 Water: To 28° 07.9' N, 177° 39.2' W.
 Vegetation: 28° 02.4' N 177° 46.3' W

Bottom: _____
 Cover: _____ Temp.: _____
 Shore: _____ Current: _____
 Dist. offshore: _____ Stream width: _____
 Depth of capture: _____ Depth of water: _____
 Collected by R. Rosenblatt + pty Styx VII
 Tide: _____ Date: Sept 21, 1968
 Method of capture: 1 KMT
 Orig. preserv.: 10% form Time: 0015 - 0828

- | | | |
|------------|--|-------------------------|
| 10 | <i>Gonostoma gracile</i> | 6 (60-113) |
| 25 | <i>Anotopterus pharao</i> | 4 (34.5-57.5) |
| 10 | <i>Chauliodus sloani</i> | 2 (48.5-51.5) |
| 65 | <i>Cryptopsaras coesii</i> | 1 (18) |
| 10 | <i>Gonostoma atlanticum</i> | 1 (60) |
| 10 | <i>Sternoptyx diaphana</i> | 5 (8-24.5) |
| 10 | <i>Argyropelecus pacificus</i> | 2 (15.5-39.2) |
| 10 | <i>Opostomias mitsuii</i> | 1 (61) |
| 25 | <i>Scopelarchus stephensi</i> | 1 (55) |
| | | HOLOTYPE |
| 25 | <i>Scopelarchus stephensi</i> | PARATYPES 2 (58.2-58.9) |
| (Cover) 8 | <i>Mentodus rostratus</i> ^{<i>triangularis</i>} _{sp #2} | 1 (13) |
| 65 | <i>Microlophichthys microlophus</i> | 1 (15) |
| 65 | <i>Edriolychnus schmidti</i> | 1 |
| 65 | <i>Dolopichthys pullatus</i> | 1 (14) |
| 65 | <i>Cryptopsaras coesii</i> | 1 (18) |
| 65 | <i>Metarocetus murrayi</i> | 1 (34) |
| 10 | <i>Diplophos taenia</i> | 1 (41.0) |
| 60 | " <i>Pseudoscopelidae</i> " [gift to USNM] | 4 (45) |
| | <i>Pseudoscopelus vorax</i> PARATYPE nsp #5 | 1 (45) |
| 65 | <i>Haplophryne mollis</i> | 1 (larva) |
| 30 | <i>Rosaera indica</i> | 1 (70.5) |
| 25 | <i>Diaphus fulgens</i> | 2 (18-57.5) |
| 26 | <i>Serrivomer</i> sp. | 1 (189) |
| 25 | <i>Myctophum nitidulum</i> | 1 (16) |
| 25 | <i>Hygophum reinhardtii</i> | 1 (31) |
| 25 | <i>Teaningichthys paurolychnus</i> | 1 (22) + 1 (61) |
| (Cover) 25 | <i>Ceratoscopus warmingi</i> | 1 (27) |

25	<i>Lampanyctus macdonaldi</i>	1(28)
25	<i>Nannobrachium achiurus</i>	1(26)
25	<i>Nannobrachium nigrum</i>	3(48-70)
25	<i>Benthoosema suborbitale</i>	4(15-27)
25	<i>Diogenichthys atlanticus</i>	1(15.5)
10	<i>Vinciguerria nimbaria</i>	2(17-19)
25	<i>Symbolophorus californiensis</i>	1(28)
10	<i>Cyclothone alba</i>	27(12-29)
10	<i>Cyclothone braueri</i>	11(18-41)
10	<i>Cyclothone acclinidens</i>	350(17-56)
10	<i>Astronesthes niger</i>	3(15- ³¹ 24)
65	" <i>Hyaloceratias</i> "	1(11)
10	<i>Astronesthidae</i> (larva)	1(29)
25	<i>Bolinichthys</i> sp	1(38)
25	<i>Lampanyctus pusillus</i>	4(20 ² -26)
25	<i>Loweina terminata</i>	1210(21-24)
25	<i>Diaphus anderseni</i>	1(16)
25	<i>Notolychnus valdiviae</i>	8(13.5- ²⁰ 18)
25	<i>Triphoturus nigrescens</i>	1(21)
25	<i>Lampadena</i> sp	1(16.5)
31	<i>Poromitra oscitans</i>	1(65)
31	<i>Scopeloberyx opisthopterus</i>	18(18.5-36)
8	<i>Tragularius mesalirus</i>	1(70)