

State or Country: Southeastern Pacific Ocean Field No. SI069-334

County: Map:, 1

Locality:

Lat. 15° 00.0' - 14° 55.05', Long. 105° 58.8' W.

Water:

Vegetation: 14° 57.5'S 106° 06.2'

Bottom:

Cover: Temp: 24.8°C

Shore: Current:

Dist. offshore: Stream width:

Depth of capture: 0-1300 m Depth of water: ca 2000 fms

Collected by R.L. Wisner & party

Tide: Date: III-21-69

Method of capture: 10' IKMT, 3500 m. W.D.

Orig. preserv.: 10% formalin Time: 0210-0815

IKMT # 4 M069-III-12

Piquero-V. Towng away from Sta. ~~77~~ 32

- | | | |
|----|--|------------|
| 31 | <i>Scopeloberyx robustus</i> | 1(80.8) |
| 10 | <i>Malacosteus niger</i> | 1(85.5) |
| 25 | <i>Cetostoma regadi</i> | 1(86) |
| 10 | <i>Gonostoma atlanticum</i> | 4(24-45.5) |
| 10 | <i>Gonostoma elongatum</i> | 3(86-200) |
| 9 | <i>Dolichopteryx longipes</i> | 1(104.7) |
| 65 | <i>Cryptopsaras couesii</i> | 1(11.5) |
| 31 | <i>Anoplogaster cornuta</i> | 2(8-16.5) |
| 26 | <i>Perichthys serpentinus</i> | 1(130) |
| 10 | <i>Valenciennellus tripunctulatus</i> | 1(19.1) |
| 25 | <i>Eurypharynx pelecanoides</i> | 1(130) |
| 31 | <i>Diracemus argenteus</i> | 1(42) |
| 10 | <i>Chauliodus sloarii</i> | 12(20-198) |
| 25 | <i>Scopelosaurus</i> sp. | 1(39.5) |
| 65 | <i>Dolopichthys longicorris</i> | 1(16.2) |
| 25 | <i>Benthalbella infans</i> | 1(61.0) |
| 65 | <i>Dolopichthys pullatus</i> | 1(16.2) |
| 25 | <i>Evermannella indica</i> | 3(21-40.4) |
| 25 | <i>Diaphus schmidti</i> | 7(26.5-41) |
| 10 | <i>Diplophos taenia</i> | 1(148.1) |
| 25 | <i>Diaphus mollis</i> | 4(20-61) |
| 31 | <i>Diracemus argenteus</i> | 1(42) |
| 25 | <i>Scopelosaurus hoedti</i> | 1(40) |
| 25 | <i>Nannobranchium gibbsi</i> - 2 specimens on EXCHANGE
to Smithsonian [USNM 303165] | |
| 25 | <i>Lampanyctus</i> n. sp.? | 4(20-29) |

9	<i>Bathylagus euryops</i>	3(20-161)
25	<i>Diogenichthys atlanticus</i>	12(11-20) 3(12-16)
25	<i>Notolychnus valdiviae</i>	26(14-21) +(20)
10	<i>Sternoptyx diaphana</i>	33(8-41)
10	<i>Idiacanthus</i> sp. <i>fasciola</i>	5 (37 1 -120)
10	<i>Argyropelecus hemigymnus</i>	2(10-22)
10	<i>Argyropelecus affinis</i>	4(18-39)
45	<i>Pentherichthys atratus</i>	1(62)
25	<i>Dmosudis lowii</i>	3(15-28)
35	<i>Bathysphyraenops simplex</i>	1(20)
25	<i>Hygophum reinhardtii</i>	4(13-41)
25	<i>Lampadena urophaos</i>	2(38.7 - 39.5)
25	<i>Lampanyctus steinbecki</i>	18(12-101)
25	<i>Lepidophanes photothorax</i>	8(16-43)
25	<i>Diaphus termophilus</i>	2(16.5-31)
25	<i>Diaphus rolfbolini</i>	3(14-23)
25	<i>Diaphus luetkeni</i>	1(40)
25	<i>Diaphus fulgens</i> (form A)	1(30.5)
25	<i>Diaphus brachycephalus</i>	4(9.5-57)
25	<i>Ceratoscopelus warmingi</i>	33(19-62)
10	<i>Ichthyococcus ovatus</i>	1(13)
25	<i>Notoscopelus resplendens</i>	1(59.5)
25	<i>Triphoturus micropterus</i>	7(18-37)
32	<i>Stethopristes eos</i>	1(24.5)
62	<i>Bregmaceros atlanticus</i>	7(24-45)
43	<i>Diplospinus multistriatus</i>	1(38)
10	<i>Vinciguerrria nimbaria</i>	36(14-38)

(continued - next page)

(continuation)

SI069-334

State or Country: Field No.

County: Map:, 1

Locality:

..... Lat. N., Long. W.

Water:

Vegetation:

Bottom:

Cover: Temp.:

Shore: Current:

Dist. offshore: Stream width:

Depth of capture: Depth of water:

Collected by

Tide: Date:

Method of capture:

Orig. preserv.: Time:

26 Serrivomeridae (lepto.) 2(71-77)