

State or Country: Southwestern Pacific Ocean Field No. SI069-348

County: \_\_\_\_\_ Map: \_\_\_\_\_, 1

Locality: \_\_\_\_\_

Lat. 17°50.0' - 41.5'S, Long. 102°39.0' W.

Water: \_\_\_\_\_

Vegetation: \_\_\_\_\_

Bottom: \_\_\_\_\_

Cover: \_\_\_\_\_ Surf. 24.500

Shore: \_\_\_\_\_ Current: 24.500

Dist. offshore: \_\_\_\_\_ Stream width: \_\_\_\_\_

Depth of capture: 0-750m Depth of water: ca 2000 fms

Collected by R.L. Wisner, et al

Tide: \_\_\_\_\_ Date: 11-8-9-1969

Method of capture: 10' Isaacs-Kidd Midwater Trawl

Orig. preserv.: 10% formalin Time: 2045-0010

Piquero - V. 1KMT # 10. MV69-III-25

10 Pachystomias microdon 1(190)

10 Chauliodus sloani 1(189)

10 Valenciennellus tripunctulatus

3(16-21)

10 Danaphos oculatus

3(31.5-34)

25 Scopelarchus aralis 3(14.8-22.6)

10 Diplophos taenia 3(31.3-45.5)

25 Diaphus sp. juv. 8(8-10)

25 Symbolophorus sp. reversus 3(20-36)

10 Argyropelecus hemigymnus 2(20-21)

10 Sternoptyx diaphana 12(8-19)

10 Idiacanthus fasciola 1(130)

25 Lampanyctus steinbecki 3(22-25)

25 Lepidophanes sp. Bolinichthys 2(12.5-18)

25 Lobjanichia gemellari 1(27.5)

25 Diaphus termophilus 1(18)

25 Diaphus schmidti 5(16-22)

25 Ceratoscopelus warmingi 1(25)

25 Benthosema suborbitale 10(16.5-28)

25 Omosudis lowii 1(17)

35 Bathysphyraenops simplex 1(9)

25 Notolychnus valdiviae 9(12-21)

26 Stemonidium hypomelas 2(93-128)

26 Serrivomer sp. 3(153-175)

25 Stemonosudis sp. 1(41)

10	Flagellostomias boureei	1(48)
31	Scopelogadus m. mizolepis	1(15)
31	Scopeloberyx opisthapterus	1(14)
62	Bregmaceros bathymaster	4(15-22)
31	Melamphaes danae	3(16-18)
10	Vinciguerria nimbaria	10(15-18)
10	Vinciguerria poweriae	4(14-24)
25	Gonichthys sp.	1(13.5)