

State or Country: CENT. NORTH PACIFIC SIO No.: SIO 70-113
County: Station: A-1

Locality: Map: _____, 1

Lat. 27° 12.1' N., Long. 155° 16.1' W.

Water: _____

Vegetation: _____

Bottom: _____

Temp.: _____ Sal.: _____ Current: _____

Dist. offshore: _____ Tide: _____

Depth of capture: 3500 mwo Depth of water: _____

Collected by Climax II exp

Time: 1835-2115

Method of capture: IKMWT 26

Orig. preserv.: Date: * Aug 1969

- | | | |
|---------------|-------------------------------|---------------------|
| 25 | Scopelarchus quentheri | 1(43.4) |
| 10 | Gonostoma atlanticum | 1(60.3) |
| 10 | Sternoptyx diaphana | 10(8-13.5) |
| 10 | Vinciguerria nimbaria | 9(10-29.5) |
| 30 | Bathyleptus lisae | 1(88.5) |
| 25 | Diaphus schmidt | 1(29.2) |
| 25 | Bolinichthys supralateralis | 1(88.5) |
| 25 | Diaphus elucens | 1(45.0) |
| 25 | Ceratoscopelus warmingi | 2(51.7-55.7) |
| 25 | Lampanyctus acanthurus | 2(53.0-53.6) |
| * 25 | Lampanyctus N. sp. | 1(50.8) |
| 25 | Symbolophorus evermanni | 1(25.3) |
| 25 | Taaningichthys bathyphilus | 1(67.7) |
| 10 | Cyclothone pallida | 29(26.8-56.0) |
| 10 | Cyclothone pseudopallida | 9(20.1-35.8) |
| 10 | Cyclothone alba | 8(19.0-26.3) |
| 25 | Scopelarchus | 1(276.8) |
| 26 | Setriovomer sp. | discarded 1(276.8) |
| 10 | Gonostoma elongatum | discarded 1(35.8) |
| 31 | Melamphaes simus | 2(26.3-26.5) |
| 31 | Poromitra oscitans | 1(53.3) |
| 31 | Scopelogadus m. mizolepis | 1(46.1) |
| 31 | Scopeloberyx opisthopterus | 2(28.8-30.5) |
| 10 | Sternoptyx diaphana | 1(11.5) |
| | Myctophid (mangled) | 1() |
| 25 | Hygophum ? proximum? | 1(24.2) |
| * 25 | Nannobranchium hawaiiensis | 2(13-50.8) |

25 Diaphus mollis #1
23 Diaphus fulgens (form A) } 2 (13.0-58.9)
— } 1 (24)