


SIO No.: SIO 70-121  
 State or Country: \_\_\_\_\_ Station: B-5  
 County: \_\_\_\_\_ Map: \_\_\_\_\_, 1  
 Locality: South Central Pacific  
 \_\_\_\_\_  
start Lat. 24°56'S ■ Long. 154°50' W  
 Water: \_\_\_\_\_  
 Vegetation: \_\_\_\_\_  
 Bottom: \_\_\_\_\_  
 Temp.: \_\_\_\_\_ Sal.: \_\_\_\_\_ Current: \_\_\_\_\_  
 Dist. offshore: \_\_\_\_\_ Tide: \_\_\_\_\_  
 Depth of capture: \_\_\_\_\_ Depth of water: \_\_\_\_\_  
 Collected by Climax II exp, Leg II  
R/V "Argo" Time: 0019-0056  
 Method of capture: IKMWPT  
 Orig. preserv.: \_\_\_\_\_ Date: 8 Oct 1969  
[Scan VII]

- |    |  |                        |
|----|--|------------------------|
| 31 | <i>Aroplogaster cornuta</i>                    | 1(9.2)                 |
| 10 | <i>Gonostoma atlanticum</i>                    | 1(27)                  |
| 10 | <i>Chauliodus sloani</i>                       | 1(109.2)               |
| 26 | <i>Derichthys serpentinus</i>                  | 1(72.5)                |
| 10 | <i>Eustomias gibbsi</i>                        | PARATYPE               |
| -  |  | PARATYPE 1(68.5)       |
| 25 | <i>Evermanella indicamegalops</i>              | 1(24.1)                |
| 25 | <i>Coccorella atrata atlantica</i>             | 1(51.9)                |
| 60 | <i>Chiasmodon cf. subniger</i>                 | 1(20)                  |
| 25 | <i>Scopelarchoides climax</i>                  | paratypes 2(20.0-35.5) |
| 25 | <i>Scopelarchoides signifer</i>                | 2(22.6-37.1)           |
| -  |  | paratypes              |
| 10 | <i>Argyropelecus aculeatus</i>                 | 1(23.5)                |
| 62 | <i>Bregmaceros</i> sp.                         | 2(17.5-21.5)           |
| 10 | <i>Gonostoma longipinnis</i>                   | 3(24.2-37.3)           |
| 10 | <i>Photostomias</i> sp.                        | 1(37)                  |
| 10 | <i>Idiacanthus fasciola</i>                    | 2(45-103)              |
|    | Melamphaid small                               | 12                     |
| 25 | <i>Triphoturus nigrescens</i>                  | 1(27)                  |
|    | Gemmylids                                      | 3                      |
| 10 | <i>Vinciguerria</i> sp. [attenuata & poweriae] | 13(6-32)               |
| 25 | <i>Ceratoscopus warmingi</i>                   | 5(15-28)               |
| 25 | <i>Diogenichthys atlanticus</i>                | 4(13-23)               |
| 25 | <i>Lampadena urophaos</i>                      | 1(37)                  |
| 25 | <i>Notoscopus resplendens</i>                  | 1(28)                  |



25	<i>Notolychnus valdiviae</i>	43(12-21)
25	<i>Symbolophorus evermanni</i>	2(22-33)
25	<i>Lobianchia</i> cf. <i>gemellarii</i>	1(18)
25	<i>Diaphus fragilis</i>	1(57)
25	<i>Taaningichthys minimus</i>	1(22)
25	<i>Lampanyctus steinbecki</i>	7 <del>6</del> (19.5-47)
25	<i>Scopelosaurus hamiltoni</i>	2(34-40)
25	<i>Benthabella infans</i>	2( <del>8</del> <sup>14</sup> -18)
25	<i>Scopelarchus analis</i>	2(8-17)