

State or Country: C.N. PACIFIC SIO No.: SIO 71-306  
 County: \_\_\_\_\_ Station: ARIES 9- H13  
 Locality: \_\_\_\_\_ Map: \_\_\_\_\_, 1  
27° 26.9' - 26.3' N  
155° 28.7' - 16.8' W  
 Lat. 27° 26.6' N., Long. 155° 22.7' W.  
 Water: \_\_\_\_\_  
 Vegetation: \_\_\_\_\_  
 Bottom: \_\_\_\_\_  
 Temp.: \_\_\_\_\_ Sal.: \_\_\_\_\_ Current: \_\_\_\_\_  
 Dist. offshore: \_\_\_\_\_ Tide: \_\_\_\_\_  
 Depth of capture: \_\_\_\_\_ Depth of water: \_\_\_\_\_  
 Collected by McGowan/Barnett Time: 0700 - 1030  
 Method of capture: 1 KMT 3000 mws  
 Orig. preserv.: formalin Date: 1 Oct. 1971

|    |                                                 |                         |
|----|-------------------------------------------------|-------------------------|
| 10 | <i>Chauliodus sloani</i>                        | 1(17.8)                 |
| 25 | <i>Omosudis lowii</i>                           | 1(23.4)                 |
| 10 | <i>Margrethia obtusirostra</i>                  | 1(18.3)                 |
| 30 | <i>Bathyleptus lisae</i>                        | 2(14.1-63)              |
| 31 | <i>Scopelogadus m. mizolepis</i>                | 3(19-19.4)              |
| 26 | <i>Serrivomer</i> sp.                           | 1(302.5)                |
| 10 | <i>Argyropelecus hemigymnus</i>                 | 2(8.3-16.2)             |
| 65 | <i>Cryptopsaras couesii</i>                     | 1(10.1)                 |
| 25 | <i>Scopelarchus quertheri</i>                   | 1(61.3)                 |
| 25 | <i>Berthosema suborbitale</i>                   | 3(11.3-31.6)            |
| 25 | <i>Ceratospilus warmingi</i>                    | 1(56.1)                 |
| 10 | <i>Vinciguerria nimbaria</i>                    | 3(15.4-29.2)            |
| 10 | <i>Sternoptyx diaphana</i>                      | 12(7.2-26)              |
| 25 | <i>Lepidopteryx longipes</i>                    | 9(17.1-46)              |
| 10 | <i>Cyclothone pseudopallida</i>                 | 14(19.8-41.3)           |
| 31 | <i>Scopeloberyx opisthopterus</i>               | 10(13.9-38.1)           |
| *  | 25 <i>Lobiarctia gemellarii</i>                 | 3(22.6-45.8)            |
|    | 31 <i>Scopeloberyx</i> n. sp.                   | 4(14.1-15)              |
|    | 62 <i>Bregmaceros japonicus</i>                 | 5(22.8-58.8)            |
|    | 10 <i>Valenciennellus tripunctulatus</i>        | 2(21-27.1)              |
|    | 25 <i>Diaphus anderseni</i> n. sp.              | 1(24.9)                 |
| ** | <del>25 <i>Lampanyctus acherus</i></del> n. sp. | <del>2(23.4-26.6)</del> |
|    | 31 <i>Scopeloberyx robustus</i>                 | 4(14.1-15)              |
|    | 10 <i>Cyclothone pallida</i>                    | 60(55-65.9)             |

|     |                                  |               |
|-----|----------------------------------|---------------|
| 10  | <i>Cyclothone alba</i>           | 62(18.3-38.2) |
| 10  | <i>Sternoptyx pseudobscura</i>   | 2(9.5-50.0)   |
| 10  | "                                | 2(9.3-25.8)   |
| 10  | <i>Sternoptyx diaphana</i>       | 2(17.0-22.0)  |
| 10  | <i>Sternoptyx indeterminate</i>  | 8(7.5-9.1)    |
| 62  | <i>Bregmaceros maclellandi</i>   | 1(30.9)       |
| 25  | <i>Diaphus rolfbolini</i>        | 5(13.6-37.8)  |
| 25  | <i>Hygophum reinhardti</i>       | 6(24.0-49.1)  |
| 25  | <i>Rondeletia loricata</i>       | 1(17.4)       |
| 25  | <i>Scopelosaurus hoedti</i>      | 2(44.8-54.4)  |
| 25  | <i>Triphoturus nigrescens</i>    | 18(15.2-34.4) |
| *25 | <i>Nannobrachium hawaiiensis</i> | 2(23.4-26.6)  |
| 25  | <i>Lestidiops Nr. Mirabilis</i>  | 1(32.7)       |